



Gramin Krishi Mausam Sewa
India Meteorological Department
Orissa University of Agriculture and Technology
RRTTS, Chiplima



Dr. Ipsita Kar
Technical Officer (I/C), GKMS

Week no-05

No.10(Eng .Bulletin)
Date:03.02.2023

West Central Table Land Zone

Sonepur

According to IMD, no rainfall is expected in the upcoming 5 days. The maximum temperature is likely to range between 30-32^oC and the minimum temperature between 11-13^oC. Clear sky to mainly cloudy sky is expected in the upcoming 5 days. The maximum relative humidity may vary between 21-33% and minimum relative humidity may vary between 8-12%. Light air is likely to blow in SSW direction on Saturday and Monday, S direction on Sunday with WSW direction on the remaining days.

Date	04/02/2023	05/02/2023	06/02/2023	07/02/2023	08/02/2023
Rainfall (mm)	0	0	0	0	0
Max Temperature (° C)	30	30	31	32	32
Min Temperature (°C)	11	12	13	13	13
Total cloud cover (okta)	1	0	0	0	0
Max Relative Humidity (%)	21	22	27	31	33
Min Relative Humidity (%)	8	8	10	12	11
Wind speed (km/h)	5	5	4	6	6
Wind direction (deg)	216	180	202	252	251

For further information, contact the IMD, Bhubaneswar, Tel. # 0674-2596116

Agromet Advisory

Rice

- For controlling Stem borer, application of Phorate @ 10 kg/acre should be done during transplanting. If the weeds are not controlled, the farmers should go for application of post-emergence application of Pyrazosulfuron @ 80g per acre, or Almix @ 8 g/acre after 15-20 days of transplanting.
- Irrigation should be provided as and when required especially at the critical stages like tillering, panicle initiation, flowering and milking stage. If paddy is 1 month old, then go for 1st top dressing with urea.

Potato

- The current weather is favorable for infestation of Cut worm in potato. Apply 1% Chloropyriphos dust or 2% Methylparathion dust over the bunds of the affected field to restrict the entry of the insects in to the unaffected field.
- In case of early blight in Potato, use copper based fungicide like Mancozeb 75 WP@ 2g/lit at 15 days interval three to four times. Destroy the diseased plant parts properly.

Sunflower

- Grow High Yielding Varieties like Modern and Hybrid Varieties PAC 36, KBSH, MSFH-8 and SFH 17. Apply 2 tons of FYM along with basal application of 8:16:12 kg NPK/acre for HYV and double the dose of fertilizer for the Hybrids. Apply Phosphorus in form of SSP to increase the oil content. Before sowing, the seeds should be treated @ 2 g Carbendazim per Kg of seeds.

Groundnut

- For controlling weeds in Groundnut, the farmers should apply Imazathapyr @ 1.5ml/lit .

Brinjal

- In case of Fruit and Shoot borer in brinjal, go for application of 60ml Coragen 20 SC in 200 lt water/acre. In brinjal wilt is seen. To control, mix 20g Streptocyclin in 200lt water and spray in the root zone.

Tomato

- In case of Fruit borer in Tomato, apply 80 ml Spinosad 14.5SC in 200 lt water/acre.

Cabbage and Cauliflower

- In case of Diamond back moth in cole crops like cabbage, cauliflower, go for application of 80 ml Spinosad 14.5SC in 200 lt water/acre.

Broccoli

- Seed rate for broccoli is 300grams/ha. Raise the seeds in the nursery bed and transplant the seedlings at 4-5 leaves or 25-30 days stage and transplant at a spacing of 40-45cm.

Chilli and Capsicum

- In case of Capsicum and chilli thrips, go for application of 200ml Dimethoate 30 EC in 200lt water/acre. In case of mites, apply 500ml Dicofol 18.5EC in 200lt water/acre.

Carrot

- Mulching in carrot should be done to protect them from direct exposure of sunlight and conserve moisture.

Technical Officer